

BUFO GUTTATUS (Guianan Golden Toad). VENEZUELA: ESTADO APURE: Santa María del Orinoco, ca. 20 km S mouth of Río Cinaruco on the Orinoco River. 21 April 2000. J. C. Paz. Museo de Biología de la Universidad del Zulia (MBLUZA-0139). Verified by Andrés Orellana. First state record. Another Amazonian species discovered NW of previous known distribution (Barrio 1998, Acta Biol. Venezuela 18[2]:1–93). The Río Cinaruco apparently constitutes a northern Amazonian reduct into the Llanos ecotone on the northwestern side of the Orinoco River (previously considered a barrier for many species, including amphibians) or is the natural limit for some Amazonian amphibians and reptiles into the Llanos. Northern Amazonian species such as the Bighead Turtle (*Peltocephalus dumerilianus*) (Pritchard and Trebbau 1984, Turtles of Venezuela. Publ. SSAR. 403 pp.), Gladiator Frog (*Hyla boans*) (Arrington and Arrington 2000, Herpetol. Rev 31:170) and Rocky Frog (*Leptodactylus lithonaetes*) (Heyer 1995, Proc. Biol. Soc. Washington 108[4]:695–716) showed similar distributions crossing the Orinoco River and reaching the Cinaruco River to the Northwest.

Submitted by **CÉSAR LUIS BARRO**, Fundación Andígena, Apartado Postal 210, Mérida 5101-A, Venezuela (e-mail: cesarlba@yahoo.com), **TITO BARROS**, Museo de Biología, Departamento de Biología, Facultad Experimental de Ciencias, La Universidad del Zulia, Apartado 526, 4001, Maracaibo, Estado Zulia, Venezuela (e-mail: trbarros@cantv.net), and **JUAN CARLOS PAZ**, Departamento de Biología, Facultad de Ciencias, La Universidad del Zulia, Maracaibo, Estado Zulia, Venezuela, Apartado Postal 526 (e-mail: j-c1971@hotmail.com).